

(Pseudo)-Dirac neutrinos and leptogenesis*

Steven Abel and Véronique Pagé

Centre for Particle Theory,

University of Durham, Durham, DH1 3LE, UK

2nd February 2008

Abstract

We discuss how Dirac neutrinos can naturally be generated in supersymmetry and how they allow for an Affleck-Dine leptogenesis scenario, in which a left-right asymmetry is generated in the sneutrino sector, the left part of which is transferred to a baryon asymmetry via sphaleron transitions. No exotic fields need to be added to the MSSM other than the right-handed neutrino.

1 Introduction

As the nature of neutrinos still eludes us, it is worth bearing in mind all the alternative possibilities for neutrino masses. In particular the Majorana or Dirac nature of neutrinos has not yet been established and, in spite of (or perhaps because of) the current hegemony of the see-saw mechanism, it is interesting to ask whether the latter scenario is a reasonable possibility. This talk is based on two papers that demonstrate the naturalness of Dirac neutrinos in the context of supergravity, paying particular attention to their masses [1] and to baryogenesis [2]. These are the two issues that need to be addressed if Dirac or pseudo-Dirac neutrinos are to be considered a viable alternative to the more usual see-saw mechanism.

Let us begin by recalling the basics; considering only one neutrino flavour for simplicity, the most general neutrino mass matrix can be written as

$$- \mathcal{L}_{mass} = \frac{1}{2} \overline{\chi}_L^c M_\nu \chi_L + \frac{1}{2} \overline{\chi}_L M_\nu \chi_L \quad (1)$$

with

$$\chi = \begin{pmatrix} \nu \\ \nu^c \end{pmatrix}, \quad \overline{\chi}^c = -(\overline{\nu^c}, \overline{\nu}),$$

$$M_\nu = \begin{pmatrix} m_L & m_D \\ m_D^T & M_N \end{pmatrix}. \quad (2)$$

*Talk given at the Dark Side of the Universe Conference 2006 (DSU2006), Madrid, Spain, 20-24 June 2006

The various scenarios for the neutrino mass are essentially differentiated by the importance of each term in the mass matrix M_ν . In the see-saw scenario [3–11], the mass terms are such that

$$M_{\nu,ss} = \begin{pmatrix} m_L & \ll & m_D \\ m_D^T & \ll & M_N \end{pmatrix} . \quad (3)$$

There is little oscillation between active and sterile because of small mixing angle. The ‘sterile’ and ‘active’ states ν^c are very nearly mass eigenstates and, when giving it a Dirac mass m_D similar to other fermions, the active state obtains naturally a very small mass m_D^2/M_N . The origin of the mass scale M_N , however, is left unexplained. At the opposite extreme, the pure Dirac neutrino has a mass matrix

$$M_{\nu,d} = \begin{pmatrix} 0 & m_D \\ m_D^T & 0 \end{pmatrix} , \quad (4)$$

and there is no oscillation between active and sterile because of mass degeneracy. Here there is no new energy scale to be explained, apart from, obviously, the neutrino mass scale itself. This has been often presented as a major drawback for the Dirac neutrino scenario as opposed to the see-saw one, however a natural explanation *does* exist in the context of supergravity, related to the solution of the μ -problem by Giudice and Masiero [12]. The μ problem is to explain why the mass term for the higgs fields $\mu H_u H_d$ should have a value similar to the supersymmetry breaking scale, to which it is apparently unrelated. The explanation of ref. [12] is that the μ -term does not appear at first order in the superpotential, but does appear in the Kähler potential of supergravity. Thus if

$$K \supset H_u H_d + H.c. , \quad (5)$$

then a term $\mu H_u H_d$ is communicated to the effective potential by gravity in the same way as the supersymmetry breaking, and hence with $\mu \sim M_{SUSY} \sim 1$ TeV.

Could such an explanation work for Dirac neutrino masses as well? The numbers certainly suggest that it could, as has been occasionally noted in the literature [13–17]. Consider for instance supergravity with Kähler potential K and superpotential W ; let us remove the neutrino mass terms from the superpotential and place them instead in the Kähler potential, in the form

$$K \supset \frac{L H_u \bar{N}}{M} + \frac{L H_d^* \bar{N}}{M} + H.c. \quad (6)$$

with $\bar{N}$ being the RH neutrino and M the scale at which higher dimensional operators make their appearance in K . Assume now that SUSY breaking is communicated by gravity, with $m_{3/2} \sim 1$ TeV, and for the sake of the argument, consider $M \approx M_P$. Then, taking $\langle H_u \rangle \approx m_{top}$, we obtain

$$m_\nu \approx \frac{m_{3/2}}{M} (\langle H_u \rangle + \langle H_d \rangle) \sim 10^{-4} \text{eV} , \quad (7)$$

which is intriguingly close to the measured value of $0.04 - 0.05 \text{eV}$, assuming that the mass-squared differences are indicative of the actual masses. A

detailed calculation has been made in [1], along with the suggestion of an R-symmetric model that could accomodate such a scenario. Here we shall simply insist on the fact that the inclusion of the neutrino mass-scale 'problem' within the framework of Susy and SUGRA allows for the appearance of scenarios that do not require the addition of a new mass scale.

2 Affleck-Dine Dirac leptogenesis

Let us now turn to the question of the leptogenesis mechanism. The original leptogenesis scenario [18] requires the presence of a Majorana mass and its accompanying mass scale. A leptogenesis scenario in the absence of a Majorana mass, sometimes called *neutrino genesis*, was introduced by Dick & al [19]. That model was presented in the context of a modified SM with an additional heavy Higgs-like doublet. The main feature was the smallness of the neutrino Yukawas which effectively hides a leptonic asymmetry away from the sphalerons, which are blind to the RH sector. Various implementations of this scenario have been suggested [20, 21]. Here we point out that, if we are willing to include the question of Dirac leptogenesis within the framework of SUSY, the Affleck-Dine (AD) mechanism [22] allows a very efficient implementation of neutrino genesis in just the MSSM + right-handed neutrinos. The AD mechanism allows the production of a $(\tilde{\nu}_L - \tilde{\nu}_R)$ current and, although lepton-number is conserved, only the LH lepton number can be converted to a baryon number through sphalerons, while the RH component is hidden by the smallness of the Yukawa coupling [2].

Let us first review the original neutrino genesis scenario and the model suggested by [19]. The model suggested requires the addition of an heavy, Higgs-like doublet. Let us call this doublet ϕ ; by the usual 'drift and decay' mechanism, the decay of ϕ starts a chain of reactions that leads to the desired baryon asymmetry:

$$\phi \rightarrow \nu_L + \bar{\nu}_R \rightarrow -\alpha(B + L) + \nu_L + \bar{\nu}_R . \quad (8)$$

Because of the smallness of the neutrino mass ν_R is inert, yet it still holds a lepton number. As usual, sphalerons transfer (only the LH) lepton number to baryons. CP violation is provided by the decay of ϕ . Our proposal is that the AD mechanism [22] can play the role of the ϕ decay, thus rendering the addition of the ϕ field unnecessary. In the AD mechanism, scalar fields 'slow-roll' along flat directions of the superpotential, which causes them to accumulate some quantum number. The original AD scenario saw the fields accumulating a B -number directly. Here they will instead accumulate a 'left-right' (LR) asymmetry, allowing neutrino genesis to create the baryon number of the Universe.

Consider the superpotential of the *effective* global Susy theory:

$$W = Y_U Q H_u U^c + Y_D Q H_d D^c + Y_E L H_d E^c + Y_\nu L H_u N^c + \mu H_u H_d . \quad (9)$$

It possesses two D -flat directions, $L H_u$ and N^c :

$$L = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi \\ 0 \end{pmatrix}, \quad H_u = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \phi \end{pmatrix}, \quad N^c = \tilde{\bar{\nu}} . \quad (10)$$

We should note that these directions are not perfectly F -flat due to the presence of the Yukawa couplings; however these are small enough not to endanger the success of the mechanism, and indeed it is the non-zero F -terms that will indirectly determine the baryon number. The scalar potential in which these fields are evolving is given by:

$$\begin{aligned} V &= V_{SB} + V_{Hubble} + V_F \\ &= (m_\phi^2 - c_\phi H^2) |\phi|^2 + (m_{\tilde{\nu}}^2 - c_{\tilde{\nu}} H^2) |\tilde{\nu}|^2 \\ &\quad + (Y_\nu (A + c_A H) \phi^2 \tilde{\nu} + h.c.) + \frac{|Y_\nu|^2}{4} |\phi|^2 + |Y_\nu|^2 |\tilde{\nu} \phi|^2 \end{aligned} \quad (11)$$

where V_{SB} , V_{Hubble} and V_F are the SUSY-breaking, 'Hubble' and F -term potentials, respectively ¹. We can see now how the LR asymmetry:

$$n_{LR} = n_L - n_R \quad (12)$$

where

$$n_L = \frac{i}{2} (\dot{\phi}^* \phi - \dot{\phi} \phi^*), \quad n_R = -i (\dot{\tilde{\nu}}^* \tilde{\nu} - \dot{\tilde{\nu}} \tilde{\nu}^*) \quad (13)$$

has a non-trivial evolution. Indeed the evolution of n_{LR} is obtained through solving the equation of motion for each field, of the type:

$$\ddot{\phi} + 3H\dot{\phi} + \frac{\partial V}{\partial \phi} \phi. \quad (14)$$

Doing this, we obtain:

$$n_{LR} + 3Hn_{LR} = 4\text{Im} (Y_\nu A \phi^2 \tilde{\nu}) . \quad (15)$$

The behaviour of the asymmetry is given in Figure (1), along with the evolution of the scalar field ϕ ; the approximate analytical behaviour for each phases of evolution is also given in Table (1). We should mention that it is obviously necessary that the $\tilde{\nu}$ oscillations decay after the electroweak phase transition. With the lifetime given by $\tau_{decay} = 4\pi/(Y_\nu^2 m_{\tilde{\nu}})$ we see that indeed $T_{decay} \sim 100\text{MeV}$. The LH sneutrinos, however, will quickly decay to neutrinos (this is instantaneous; see [2]), and this brings us back to original neutrino genesis. Sphalerons can transfer this (LH) lepton number to baryons, although at all times the LH lepton number is accompanied by an equal and opposite lepton number hidden (to sphalerons) in the inert RH sneutrinos. The fact that the RH, sterile sneutrinos hold an asymmetry equal and opposite to the baryonic one opens the interesting possibility that they form the cold component of dark matter.

¹The 'Hubble' potential is the effective potential due to SUSY breaking in the early Universe. It is the presence of these terms that allow the flat directions to develop large expectation values during inflation - see [23, 24]

$H \gg m_{3/2}$	$R \sim t^{2/3}$	chaotic motion	$R^3 H^2 \phi^2 = \text{const} \Rightarrow n_{LR} \sim \text{const}$
$H(T_R) < H < m_{3/2}$	$R \sim t^{2/3}$	cyclic motion	$R^3 m_\phi^2 \phi^2 = \text{const} \Rightarrow n_{LR} \sim t^{-1}$
$H < H(T_R)$	$R \sim t^{1/2}$	cyclic motion	$R^3 m_\phi^2 \phi^2 = \text{const} \Rightarrow n_{LR} \sim t^{-3/2}$

Table 1: Approximate analytical behaviour of the LR asymmetry. The early chaotic period is typical of the flat directions used here, as those are not perfectly F -flat.




Figure 1: Time evolution of the generated LR asymmetry for typical parameters. The behaviour of the ϕ field is also shown for early (shortly before $H \sim 100 \text{ GeV}$) and late (post-reheating) times.

3 The baryonic asymmetry

To be able to approximate the LR asymmetry, let us recall that before reheating, while the inflaton oscillations dominate the energy of the Universe, both the inflaton and the asymmetry behave like matter. Thus we can use the inflaton energy density to keep track of the asymmetry:

$$\frac{\rho_{n_{LR}}}{\rho_I} \sim \frac{m_{3/2}^2 |A/Y_\nu|^2}{m_{3/2}^2 M_P^2} . \quad (16)$$

After reheating, it is the ratio of the asymmetry with entropy that remains constant, and now $\rho_{n_{LR}} = m_{\phi, \tilde{\nu}} n_{LR}$. Hence the asymmetry is:

$$n_{LR} \approx \frac{|A/Y_\nu|^2}{M_P^2} \frac{T_R}{m_\phi} = 10^{-9} \left| \frac{A}{100 \text{ GeV}} \right|^2 \left| \frac{10^{-12}}{Y_\nu} \right|^2 \left| \frac{T_R}{1 \text{ TeV}} \right| \left| \frac{100 \text{ GeV}}{m_\phi} \right| \quad (17)$$

This asymmetry is related to the baryon number in a fairly straightforward way [25, 26]:

$$B = L = \frac{8}{23} n_L^{(R)} \quad T > T_{ew} , \quad (18)$$

Hence the observed baryon number is obtained for reheating temperatures of order 1 TeV. Should the RH sneutrinos be the dark matter, this last relation allows us to constrain their mass via

$$m_{DM} = \frac{8}{23} \frac{\Omega_{DM}}{\Omega_b} m_b . \quad (19)$$

4 Conclusion

The possibility of neutrinos being Dirac or pseudo-Dirac particles is still very much alive. First, within the framework of supergravity, neutrino masses can be made naturally small. Their scale can be related to the scale hierarchy between the weak and Planck scale, in much the same way as for the μ -term. No new scale is necessary to explain the neutrino mass. Moreover, still within SUSY, leptogenesis with Dirac neutrinos can be easily implemented using the Affleck-Dine mechanism. With a reheating temperature of order 1TeV, the right order of magnitude for the baryon number of the Universe is obtained. This requires no new fields to be added to the MSSM beyond the right handed neutrino. In the case of pure Dirac neutrinos the $B - L$ number of the visible sector is connected to an equal and opposite right-handed sneutrino number, and this provides an intriguing connection between the dark matter density and the baryon number of the Universe. This link has been obtained in the past in various works [27–37], but here we see it arising quite naturally.

References

- [1] S. A. Abel, A. Dedes, and K. Tamvakis. *Phys. Rev.*, D71:033003, 2005.
- [2] Steven Abel and Veronique Page. *JHEP*, 05:024, 2006.
- [3] Peter Minkowski. *Phys. Lett.*, B67:421, 1977.
- [4] M. Gell-Mann, M.P Ramond, and R. Slansky. In P. Van Nieuwenhuizen and D. Z. Freedman, editors, *SUPERGRAVITY. PROCEEDINGS, WORKSHOP AT STONY BROOK, 27-29 SEPTEMBER 1979*. Amsterdam, Netherlands: North-holland (1979) 341p.
- [5] In O. Sawada and A. Sugamoto, editors, *PROCEEDINGS OF THE WORKSHOP ON THE UNIFIED THEORY AND THE BARYON NUMBER IN THE UNIVERSE*. Tsukuba, Japan: Natl.Lab.High Energy Phys.(1979) 109 P.and Japan Natl Lab High Energy - KEK-79-18 (79,REC.JAN 80) 109p.
- [6] R. N. Mohapatra and G. Senjanovic. *Phys. Rev. Lett.*, 44:912, 1980.
- [7] Steven Weinberg. *Phys. Rev.*, D22:1694, 1980.
- [8] J. Schechter and J. W. F. Valle. *Phys. Rev.*, D22:2227, 1980.
- [9] G. Lazarides, Q. Shafi, and C. Wetterich. *Nucl. Phys.*, B181:287, 1981.
- [10] J. Schechter and J. W. F. Valle. *Phys. Rev.*, D25:774, 1982.
- [11] J. W. F. Valle. 2006.
- [12] G. F. Giudice and A. Masiero. *Phys. Lett.*, B206:480–484, 1988.
- [13] N. Arkani-Hamed, L. J. Hall, H. Murayama, D. R. Smith, and N. Weiner. *Phys. Rev.*, D64:115011, 2001.
- [14] F. Borzumati and Y. Nomura. *Phys. Rev.*, D64:053005, 2001.
- [15] J. A. Casas, J. R. Espinosa, and I. Navarro. *Phys. Rev. Lett.*, 89:161801, 2002.

- [16] R. Kitano. *Phys. Lett.*, B539:102–106, 2002.
- [17] R. Arnowitt, B. Dutta, and B. Hu. *Nucl. Phys.*, B682:347–366, 2004.
- [18] M. Fukugita and T. Yanagida. *Phys. Lett.*, B174:45, 1986.
- [19] K. Dick, M. Lindner, M. Ratz, and D. Wright. *Phys. Rev. Lett.*, 84:4039–4042, 2000.
- [20] D. G. Cerdeno, A. Dedes, and T. E. J. Underwood. 2006. hep-ph/0607157.
- [21] Brooks Thomas and Manuel Toharia. 2006. hep-ph/0607285.
- [22] I. Affleck and M. Dine. *Nucl. Phys. B*, 249:361–380, 1985.
- [23] M. Dine, L. Randall, and S. Thomas. *Nucl. Phys.*, B458:291–326, 1996.
- [24] M. Dine, L. Randall, and S. Thomas. *Phys. Rev. Lett.*, 75:398–401, 1995.
- [25] J.A Harvey and M.S Turner. *Phys. Rev. D*, 42(10), 1990.
- [26] H. K. Dreiner and G. G. Ross. *Nucl. Phys.*, B410:188–216, 1993.
- [27] S. M. Barr, R. S. Chivukula, and E. Farhi. *Phys. Lett.*, B241:387–391, 1990.
- [28] S. M. Barr. *Phys. Rev.*, D44:3062–3066, 1991.
- [29] D. B. Kaplan. *Phys. Rev. Lett.*, 68:741–743, 1992.
- [30] S. D. Thomas. *Phys. Lett.*, B356:256–263, 1995.
- [31] V. A. Kuzmin. *Phys. Part. Nucl.*, 29:257–265, 1998.
- [32] A. Kusenko. 1998. hep-ph/9901353.
- [33] G. R. Farrar and G. Zaharijas. 2004. hep-ph/0406281.
- [34] D. Hooper, J. March-Russell, and S. M. West. *Phys. Lett.*, B605:228–236, 2005.
- [35] R. Kitano and I. Low. *Phys. Rev.*, D71:023510, 2005.
- [36] N. Cosme, L. Lopez Honorez, and M. H. G. Tytgat. *Phys. Rev.*, D72:043505, 2005.
- [37] R. Kitano and I. Low. 2005. hep-ph/0503112.